

PROFESSIONAL INFORMATION

SCHEDULING STATUS

S0

1. NAME OF THE MEDICINE

Vitatech Testo Support Tablets 30

2. COMPOSITION

Each tablet contains:

Ingredients	Elemental	NRV% *
Vitamin B ₆ (as Pyridoxine Hydrochloride)	1,7 mg	100%
Vitamin D ₃ (as dl-Alpha Tocopherol Acetate)	15,0 mcg	100%
Magnesium (as Oxide)	129,0 mg	30%
Zinc (as Gluconate)	11,0 mg	100%
Boron (as citrate)	100,0 mg	-

* South African Nutrient reference values for individuals 4 years and older

For full list of excipients, see section 6.a

3. PHARMACEUTICAL FORM

White, round or oval tablet.

4. CLINICAL PARTICULARS

(a) Therapeutic indication

The vitamins and minerals in **Vitatech Testo Support** assist to maintain optimal nutritional support and wellness in the form of good health through a biochemical process at cellular level.

(b) Dosage and method of administration

Adults: Take 1 tablet per day with a glass of water after a meal.

Do not exceed the recommended dose.

Always consult your healthcare professional if you are taking any other medicine.

Only take the medication as indicated on the packet or container.

If you skip a dose, do not take a double dose.

Take at least two hours before or after taking other medication.

(c) Contraindications

Hypersensitivity to any of the ingredients or the excipients. Not for use in children under the age of 18 years.

(d) Special warnings and precautions for use

Vitatech Testo Support should be used with care if patients have any of these health problems:

- (1) High calcium levels or renal disease,

- (2) If you have certain types of anaemia like pernicious anaemia or haemolytic anaemia,
- (3) If you low levels of dopamine,
- (4) If you have hypothyroidism,
- (5) If you take heart medication,
- (6) If you have gastrointestinal diseases such as IBS (irritable bowel disease), diverticulum and ulcers,
- (7) If you have convulsions or seizures,
- (8) If you have iron related conditions like Hemochromatosis or haemoglobin,
- (9) kidney disease,
- (10) heart disease,
- (11) nausea, vomiting; a blockage in your intestines;
- (12) with estrogen-sensitive conditions it should be avoided, but with it should be used cautiously with concomitant estrogens,
- (13) low levels of calcium in your blood; or
- (14) a sudden change in bowel habits for 2 weeks or longer.

Any other supplementation consisting of vitamins and minerals should be taken into account when prescribing / suggesting **VITATECH TESTO SUPPORT**.

(e) Interaction with other medicines and other forms of interaction

No interaction studies have been performed on **VITATECH TESTO SUPPORT**, however the below information gathered from the literature on the active ingredients should be taken into consideration:

Vitatech Testo Support may interact with the following medicines. Caution is advised when used concurrently:

Vitamin D₃:

with heart conditions and medication, may contribute to irregular heartbeat and a reduce response to calcium channel blockers with raised calcium levels and/or iron with high blood pressure or hypertension medication may contribute to hypercalcemia and milk-alkali syndrome if used with dairy products.

Vitamin B₆ (Pyridoxine):

seizures and epilepsy medications when used in conjunction with pyridoxine and/or folic acid may experience reduce levels of absorption.

Magnesium:

kidney disease; heart disease; nausea, vomiting; a blockage in your intestines; low levels of calcium in your blood; or a sudden change in bowel habits for 2 weeks or longer.

Zinc:

an allergy to zinc; or any condition that makes it hard for your body to absorb nutrients from food (malabsorption).

Boron: no known.

The use of other VitaTech products may reduce the absorption of penicillamine. In conjunction with any thyroid treatment:

Anti-acids, such as altretamine, cisplatin, should be taken at least 2 hours before or after **VITATECH TESTO SUPPORT**, as they may reduce the absorption of vitamins and minerals.

- certain antibiotics (e.g., chloramphenicol),
- other vitamin/nutritional supplements

- Anticoagulants/Antiplatelet medicines, like warfarin,
- may have estrogenic effects and may therefore interfere with hormone replacement therapy,
- Immunosuppressant medicines that may possess immune stimulating properties.
- certain anti-seizure drugs (e.g., phenytoin),
- an antibiotic;
- a diuretic or "water pill";
- penicillamine;
- a blood thinner--warfarin, Coumadin, Jantoven; or
- medicine to treat osteoporosis or Paget's disease--alendronate, ibandronate, risedronate, Fosamax, Boniva, Actonel, and others.

(f) Pregnant or Breastfeeding

Intravaginal boric acid has been associated with birth defects in the first 4 months of pregnancy. If you are pregnant or breastfeeding, before using **VITATECH TESTO SUPPORT**, please consult your health care provider.

(g) Ability to drive and use machines

VITATECH TESTO SUPPORT should not affect the ability of individuals to drive or operate machinery.

(h) Side Effects

Signs of an allergic reaction like:

- rash;
- hives;
- itching;
- red, swollen, blistered, or peeling skin with or without fever;
- wheezing;
- tightness in the chest or throat;
- trouble breathing, swallowing, or talking;
- unusual hoarseness;
- or swelling of the mouth, face, lips, tongue, or throat,
- upset stomach or throwing up,
- decreased sensation to touch, temperature, and vibration,
- loss of balance or coordination,
- numbness in your feet or around your mouth,
- clumsiness in your hands; or
- feeling tired.

Not all side effects reported for this medicine are included in this leaflet. Should your general health worsen or if you experience any side-effects while taking this medicine, please consult your health care provider for advice.

(i) Overdose

In the event of over-dosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

(j) Effects when no further used

There should be no side effects or any withdrawal symptoms after you stop using **VITATECH TESTO SUPPORT**

(5) PHARMACOLOGICAL PROPERTIES

(a) PHARMACOLOGICAL CLASSIFICATION

34.12 Category D (Health Supplement) [S0]

(b) PHARMACOLOGICAL ACTION

- Essential nutrients are substances required by the body to perform its basic functions and must be obtained from our diet since the human body cannot produce them. They have one or more of three basic functions: to provide energy, contribute to body structure, and/or regulate chemical processes in the body.
- They're essential for many body functions, including building strong bones and teeth, regulating your metabolism, and staying properly hydrated. These basic functions allow us to detect and respond to environmental surroundings, move, excrete wastes, breathe, grow, and reproduction.
- When essential nutrients are in short supply or is not able to be utilized properly, a specific deficiency syndrome may be the result.
- When the deficient nutrient is resupplied before irreversible damage occurs, the signs and symptoms are reversed. Deficiency of essential nutrients is noticeable in several symptoms that can be easily observed and includes, but are not limited to, for example: weakness, tiredness, decrease in energy which can also lead to the risk of developing illnesses and health problems.
- Calcium and Vitamin D contribute to the maintenance of normal muscle function and help in the development of healthy bones and teeth.
- Copper, Manganese, Vitamin C, Vitamin E and Zinc are antioxidants which contribute to the protection of cells from oxidative stress and to normal function of the immune system.
- Magnesium contributes to normal electrolyte balance, muscle function, psychological function, and the reduction of tiredness and fatigue. Magnesium oxide is used as a supplement to maintain adequate magnesium in the body and is also used as an antacid to treat indigestion, or as a laxative to relieve occasional constipation.
- Boron is a factor in the maintenance of good health. Boron has been included in nutritional supplements or natural remedies designed to improve bone and joint health. It is a beneficial bioactive element.

(6) PHARMACEUTICAL PARTICULARS

(a) Other ingredients

Microcrystalline cellulose, di-calcium phosphate & magnesium stearate.

(b) Shelf life

2 years

(c) Storage conditions

Store at or below 25 °C.

Protect from sunlight and
moisture.

Keep out of reach of children

Store in original packaging

Do not store in a fridge

Always seal the container properly before storage



(d) Presentation and packaging

White with green lid packaged container with tamperproof induction seal
Clearly labelled with detailed information printed on the label.
Containing 30 tablets.

(e) Disposal and handling

Wrap the contents and the container properly in put in a plastic bag.
Dispose the wrapped package in the normal household waste bag after dissolving it in water.

(7) NAME AND ADDRESS OF REGISTRATION HOLDER

Serfie Imports & Exports (PTY) Ltd
Cnr Parkin & Delson Street, North End, Gqeberha 6001
Tel no: +27 (0) 41 816 3100
Email: info@vitatechhealth.com
Website: www.vitatechhealth.com

(8) REGISTRATION NUMBER

To be allocated

(9) DATE OF PUBLICATION OF PROFESSIONAL INFORMATION

July 2021

(10) COMPLEMENTARY MEDICINE

This medicine has not been evaluated by the SAHPRA for its quality, safety or intended use. This medicine is not intended to diagnose, treat, cure or prevent any disease.